

<https://helda.helsinki.fi>

Genetic growth potential, rather than phenotypic size, predicts migration phenotype in Atlantic salmon

Debes, Paul V.

2020-07-29

Debes , P V , Piavchenko , N , Erkinaro , J & Primmer , C R 2020 , ' Genetic growth potential, rather than phenotypic size, predicts migration phenotype in Atlantic salmon ' , Proceedings of the Royal Society B. Biological Sciences , vol. 287 , no. 1931 , 20200867 . <https://doi.org/10.1098/rspb.2020.0867>

<http://hdl.handle.net/10138/318377>

<https://doi.org/10.1098/rspb.2020.0867>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

**Genetic growth potential, rather than phenotypic size, predicts migration
phenotype in Atlantic salmon.**

Author list: Paul V Debes^{1,2*}, Nikolai Piavchenko¹, Jaakko Erkinaro³, and Craig R Primmer^{1,4}

Correspondence: *debes@holar.is

Affiliations: ¹Organismal & Evolutionary Biology Research Program, University of Helsinki, Helsinki, 00014, Finland; ²Department of Aquaculture and Fish Biology, Hólar University College, Sauðárkrókur, 551, Iceland; ³Natural Resources Institute Finland (Luke), Oulu, 90014, Finland; ⁴Institute of Biotechnology, University of Helsinki, Helsinki, 00014, Finland

ORCIDs:

Paul V Debes: <https://orcid.org/0000-0003-4491-9564>

Jaakko Erkinaro: <https://orcid.org/0000-0002-7843-0364>

Craig R Primmer: <https://orcid.org/0000-0002-3687-8435>

Running head: Determinants of migration phenotypes

Keywords: Partial migration, age-specific migration, threshold model, Atlantic salmon, smolting

18 **Abstract**

19 Knowledge of the relative importance of genetic vs. environmental determinants of major developmental
20 transitions is pertinent to understanding phenotypic evolution. In salmonid fishes, a major developmental
21 transition enables a risky seaward migration that provides access to feed resources. In Atlantic salmon,
22 initiation of the migrant phenotype, and thus age of migrants, is presumably controlled via thresholds of
23 a quantitative liability, approximated by body size expressed long before the migration. However, how
24 well size approximates liability, both genetically and environmentally, remains uncertain. We studied 32
25 Atlantic salmon families in two temperatures and feeding regimes (fully fed, temporarily restricted) to
26 completion of migration status at age 1 year. We detected a lower migrant probability in the cold (0.42)
27 than the warm environment (0.76), but no effects of male maturation status or feed restriction. In
28 contrast, body length in late summer predicted migrant probability and its control reduced migrant
29 probability heritability by 50-70%. Furthermore, migrant probability and length showed high heritabilities
30 and between-environment genetic correlations, and were phenotypically highly correlated with stronger
31 genetic than environmental contributions. Altogether, quantitative estimates for the genetic and
32 environmental effects predicting the migrant phenotype indicate, for a given temperature, a larger
33 importance of genetic than environmental size effects.

34 1. Introduction

35 Genetic vs. environmental determinants of developmental transitions between life stages are a major
36 topic in ecological and evolutionary studies [1-3]. Generally, both the environment and genes are assumed
37 to underlie the expression of plastic developmental phenotypes [3], but the relative contribution of each
38 often remains unknown [4, 5]. However, knowledge of environmental vs. genetic contributions underlying
39 life-stage transitions is pivotal to making accurate ecological or evolutionary predictions under
40 environmental change, such as global warming [5, 6]. A major life-stage transition in many species is
41 associated with feeding or reproduction migrations, whereby the migrant phenotype, as opposed to the
42 resident phenotype [defined as individuals not migrating during a particular season; 7], may express in
43 only part of the population, or vary with age [8, 9]. Even though such partial migration, or variation of
44 migrant age, is assumed to have considerable ecological and evolutionary consequences [8-10], evidence
45 for major underlying genetic effects is scarce or controversial [11-13]. Nonetheless, selection and crossing
46 experiments in birds and fishes indicate a heritable genetic basis [8, 14], but for fishes, underlying
47 mechanisms remain elusive [reviewed in 14]. To disentangle genetic from environmental components of
48 binary traits, including migration phenotypes, the threshold model may be appropriate [10, 15]. The
49 threshold model assumes that the expression of categorical phenotypes underlies a (usually unobserved)
50 continuous liability [16-18]. Only if the liability, influenced by both environmental and genetic effects,
51 exceeds one (or several) threshold(s) during a sensitive period, the developmental transition towards the
52 alternative phenotype(s) is initiated.

53 In many fish species also the age of the developmental transition towards the migratory phenotype shows
54 considerable variation. In salmonid species, such as salmon, trout, charr, and whitefish, the age when the
55 freshwater-hatched fish migrate to the sea (or larger water bodies, such as lakes) varies considerably,
56 encompassing one to eight years in Atlantic salmon [reviewed by 19]. Importantly, within a population
57 older migrants are usually larger than younger migrants, whereby body size increases salinity tolerance
58 and predator avoidance ability [4, 19, 20]. As a result of the latter, size at seaward migration, both within
59 and across migrant ages, often covaries with migration survival [4, 21-23].

60 Whether an Atlantic salmon of a given age undergoes the required developmental transition to become
61 a migrant ("smolt") in the following spring has long been suggested to be determined by body length in
62 the previous summer or autumn [reviewed in 19]. Therefore, and because the liability remains unknown,
63 body length is often used as an *a priori* liability proxy, and many previous indirect estimates for genetic
64 migration phenotype variation stem from investigations on such proxies [reviewed in 24]. However, the
65 liability proxy-trait variation as relevant to initiating the developmental transition is expressed well in
66 advance of the actual migration [25], as is suspected for many other developmental transitions [26]. Such
67 time lag poses logistical challenges not only for organisms after they have adopted a developmental

68 transition, but also for researchers studying a dynamic liability proxy. It is thus unsurprising that - despite
69 considerable knowledge on the physiological changes and underlying mechanisms *during* the transition
70 [27] - exact mechanism *initiating* the transition from the resident towards the migrant phenotype remain
71 unknown.

72 Even though body size may be a useful proxy determining whether the migrant phenotype is initiated, it
73 may provide an inappropriate liability proxy in many cases. Specifically, initiation of the migrant
74 phenotype triggers additional growth differences between prospective migrants and residents (which
75 may migrate at older age in Atlantic salmon) that increase their size differences, thereby blurring cause
76 and effect. During the transition, prospective migrants express accelerated growth relative to residents,
77 which results in size bimodality and culminates in migrants being larger than residents at the time of
78 migration, both among and within families [25, 28-32]. Similar confounding between causes for and
79 effects of the transition towards the migrant phenotype apply to inferences about body condition.
80 Probably owing to these methodological challenges, past inferences about effects on migration
81 phenotypes were often based on size records from any time between presumed growth acceleration and
82 the time of migration [reviewed in 24], and thus statistical associations between size and migrant
83 probability are inconsistent across studies .

84 Here, we overcome a number of the abovementioned limitations and use Atlantic salmon as a model to
85 investigate genetic and environmental determinants for the probability to exhibit the migrant phenotype
86 in the second spring (hereafter called migrant probability; MIG), focussing on body size at the end of the
87 first summer as a liability proxy. Specifically, we combined longitudinal common-garden experimentation
88 in two water temperatures and two feeding regimes with quantitative genetic methodology. We followed
89 development of 663 individuals from 32 pedigreed Atlantic salmon half-sib families until their second
90 spring, when we assessed migration phenotypes (**figure 1**). We used uni- and bivariate models to test for
91 environmental effects of water temperature, feeding regime, body condition, and male maturation on
92 MIG. Further, we tested within and between two environments with a 2° C seasonal temperature
93 difference, whether body length covaried with MIG and partitioned this covariance into environmental
94 and genetic components. Our results support a long-suspected strong genetic joint-governance of body
95 size in late summer with migration phenotype in spring, which we demonstrate to be extremely stable
96 between two temperature environments. These results widen our understanding of genetic and
97 environmental importance in developmental transitions, as well as their thermal stability, thereby
98 providing quantitative primers for future modelling relying on environmental vs. genetic relationships
99 among life-history traits across environmental temperatures [e.g., 33].

1002. Material and methods

101 (a) Experimental population and settings, and data collection points and routines

102 In October 2017, we created 32 half-sib families (generation 2) by crossing 32 parents (generation 1) in
 103 eight 2x2 factorials of unrelated individuals (**figure 1 a**). The parents originated from a broodstock
 104 (generation 0) that was hatchery raised, maintained by the Natural Resources Institute Finland,
 105 Taivalkoski, Finland, but whose individuals successfully completed a sea migration [Oulu River; described
 106 in 34]. We created and reared the experimental cohort commonly in a newly established laboratory with
 107 two similar water recirculation systems at the University of Helsinki, Finland, controlled for water
 108 temperature, oxygen, dissolved nitrate components, and light. We incubated eggs and larvae in vertical
 109 incubators with two replicates per family and temperature environment. At first feeding, we pooled same-
 110 temperature family replicates and randomised an equal number of individuals from each family (3-7) to
 111 each of eight similar 250 L tank replicates per temperature environment (i.e., all families in all tanks). The
 112 fish were part of other studies, of which one required lethal sampling [35], reducing fish numbers
 113 continuously. After sampling and natural mortality, 663 individuals yielded data for migration status in
 114 May 2019 (393 in cold and 270 in the warm environment, respectively). Water temperature in egg
 115 incubators and tanks was controlled to follow a seasonal cycle, approximating a 2 °C difference referred
 116 to hereafter as “warm” and “cold” temperature treatments (warm, range: 6.3-17.7 °C; cold, range: 4.1-
 117 16.0 °C; **figure 1 b**). Tank illumination, using fluorescence lights (4000 K, 35 W; 500 Lux at the surface) and
 118 controlled by a digital astronomical time switch (without dimming), followed the natural cycle at 61.054°
 119 (latitude) and 25.042° (longitude) (**figure 1 c**).

120 Fish were fed a commercial salmon diet *ad libitum* with particle sizes meeting fish-size compositions at all
 121 times (start dates: **figure 1 b**). In August 2018 (warm) and September 2018 (cold) we anesthetized
 122 individuals (using methanesulfonate), inserted passive integrated transponder tags (into the body cavity;
 123 12 mm) to enable re-identification, and fin clipped individuals to allow for genotyping. After fasting fish
 124 for 24 h, we measured fork length (± 1 mm) between August and September 2018 - around the time of
 125 daylengths suspected to characterise the sensitive period when migration status initiation is determined
 126 [**figure 1 c**, 15-12 h; 30]. These measurements were followed by a feed-restriction-treatment period
 127 (either *ad libitum* feeding for seven days per week, or *ad libitum* feeding for two days per week with no
 128 feeding for two or three days between feedings) that was crossed with the temperature treatment for a
 129 five-week period per tank (September - October 2018; **figure 1 b, c**). We again fasted and re-measured
 130 fish after the feed-restriction treatment in October. We determined maturation status in December 2018
 131 (during the natural spawning period), when we categorised males as mature when observing milt during
 132 gently pressing the abdomen (females rarely mature at this age). We determined migration status in May
 133 2019 [during the population-specific period of highest migration success; 34] (**figure 1 b**), when we

categorised individuals as first-year migrants when showing darkened fin edges, lack of colour patches (“parr marks”) and enhanced body silvering (**figure 1 d**). We categorized individuals as residents when not displaying signs of the migrant phenotype at this time. It should be noted that many Atlantic salmon remaining resident at 1 year of age may become migrants in later years [4], and therefore the resident category also includes potential later year (older) migrants. However, it is also possible that residents, especially males, can reproduce without migrating [4, 36]. Animal experimentation followed European Union Directive 2010/63/EU under license ESAVI-2778-2018 by the Animal Experiment Board in Finland (ELLA).

(b) Pedigree construction

We determined genotypes of parents and experimental individuals using a multiplex-PCR for 177 single nucleotide polymorphisms (SNPs) of an established panel [37], and by sequencing using an Ion Torrent (984 broodstock individuals from which we drew parental individuals) or Illumina platform (MiSeq or Next-Seq) (parental individuals, experimental individuals). Using unlinked, polymorphic SNPs, we reconstructed grandparents of the experimental individuals (with 131 usable SNPs) under maximum likelihood [38], which we combined with knowledge about the crossing scheme used to create the parental generation, and assigned the experimental individuals to their 32 parents (with 141 usable SNPs) with a likelihood approach [39], resulting in a three-generation pedigree (**figure 1 a**). On this pedigree, we based the inverse relationship matrix (A^{-1}) utilized to infer additive genetic variance via animal-model analyses [40] described below.

(c) Statistical analyses

To test model terms and estimate means and (co)variance components for MIG and LEN, we used uni- and bivariate generalized animal models with probit-link function for MIG, corresponding to animal threshold models [41]. The models were fitted with the R package MCMCglmm [42] using Bayesian Markov chain Monte Carlo simulations, which appear an appropriate method for animal models on categorical data [41, 43]. Initially, we fitted a univariate linear model:

$$y = \mu + \text{Feed} + \text{Temp} + \text{Feed:Temp} + \text{Mat} + \text{Feed:Mat} + \text{Dam} + \text{Animal} + \text{Tank} + \text{Residual},$$

where y is the liability for migrant probability, μ a model constant, Feed the fixed feed restriction environment effect (full, restricted), Temp the fixed temperature restriction environment effect (cold, warm), Mat the fixed male maturation effect (0, 1), Feed:Mat the fixed interaction effect of the feed restriction with the maturation effect (restricted to the warm environment), Dam the random dam effects ($n = 32$), Animal the random additive genetic effects (736 entries in A^{-1}), Tank the random tank effects ($n = 16$), Residual the random residual effects ($n = 663$). We also extended a reduced version of the univariate model (see results) by including body length in late summer as a continuous covariate (“length”, a

predictor; log-transformed, mean centred, and variance scaled), which we interacted with the temperature treatment. We also extended the reduced univariate models to test whether MIG differed among genotypes of a locus (*vgll3*) that has been shown to have a strong effect on male maturation probability in this experimental population [35], but we did not detect *vgll3* effects, either when controlling or not controlling for phenotypic length in late summer (electronic supplementary material, **table S1**). We estimated all variances conditional on temperature environments with 2x2 covariance matrices between environments for dam (D) and animal (G) effects and diagonal covariance matrices for tank (C) and residual (R) effects. We fitted bivariate models by extending the reduced model (without the length covariate) with body length in late summer as a second response ("LEN", "length" as a response). For the bivariate model, we additionally allowed for between-trait covariances.

We ran univariate and bivariate models for 1,500,000 and 6,000,000 iterations, respectively, and sampled every 100 iterations. For each model, we ran four chains and determined i) whether the MCMC sampling had converged as indicated by a scale reduction factor around 1 per chain [44], ii) the required number of samples to discard ("burnin") until consistently reaching a scale reduction factor < 1.1 across chains [44], iii) required thinning to have autocorrelations at lag 2 < 0.1 per chain, and confirmed whether MCMC resulted in sufficient mixing using visual examination of trace plots per chain. These criteria resulted in subsequently combined posteriors across chains with sample sizes between 8,000 and 22,000. We conducted a prior sensitivity analysis as reported in the electronic supplementary material (**figures S2-S4**). We tested feed-restriction-treatment effects on growth rate by a general animal model and predicted the sexual maturation rate using a generalised animal model (electronic supplementary material, **table S2, figure S1 and table S3**, respectively)

(d) Derived parameters

We calculated heritability as the proportion of additive genetic variance (V_A) to the total phenotypic variance (V_P). Under the threshold-model interpretation of generalized model estimates that include common environmental variance (V_C), residual variance fixed to 1 (V_R), and a variance of 1 for the probit link function [45], this results for MIG in:

$$h^2 = \frac{V_A}{V_P} = \frac{V_A}{V_A + V_C + V_R + 1}.$$

We estimated correlations at the phenotypic (R_P), residual environmental (R_E), and genetic levels (R_G) as in [46], but additionally accounted for common environmental effects on R_P . To translate liability estimates to the proportional scale, we used either the "predict.MCMCglmm" function of the MCMCglmm R-package or (for heritabilities) the "QGmvparams" function of the QGglmm R-package [45]. We present mean estimates with credible intervals (95% highest posterior density estimates of model posteriors).

199 **3. Results**

200 **(a) Migrant probability is not affected by a temporary feed restriction, male maturation, or maternal**
201 **effects**

202 Using a univariate generalized animal model and not controlling for phenotypic body length, we detected
203 that MIG was higher in the 2 °C warmer environment (**table 1**). However, although the feed restriction
204 reduced the specific growth rate by about 50% (electronic supplementary material, **table S2, figure S1**),
205 MIG was not affected by this restriction (**table 1**). MIG was also not affected by male sexual maturation,
206 which occurred only in males in the warm environment at a rate of 0.19 (95% CI: 0.08-0.33; electronic
207 supplementary material, **table S3**), nor by any interaction term (**table 1**). We also did not detect any
208 maternal (dam) or common environmental (tank) effect variance on MIG (**table 1**). We therefore removed
209 dam, but not tank effects, from models because the latter constitute the experimental replicates for the
210 temperature environments.

211 **(b) Migrant probability, but not its heritability, differs between temperature environments**

212 After removing model effects not different from zero, we estimated an average MIG of 0.41 (0.27-0.55)
213 in the cold and 0.76 (0.63-0.87) in the warm environment (probit scale contrast: 1.68, 0.95-2.36). We also
214 estimated relatively high heritabilities (h^2) for MIG liability ($h^2_{Cold} = 0.53, 0.33-0.72$; $h^2_{Warm} = 0.43, 0.22-$
215 0.64), that were not different between temperature environments (thereby rejecting evidence for one
216 component of genotype-by-environment interactions; $h^2_{Cold} - h^2_{Warm} = 0.10, -0.17-0.36$). We further
217 estimated a high between-environment genetic correlation ($R_{G_{Cold,Warm}}$) with the upper 95% credible
218 interval being very close to unity ($R_{G_{Cold,Warm}} = 0.84, 0.62-0.99$), indicating that genotype re-ranking for MIG
219 was negligible between the temperature environments (thereby rejected another aspect of genotype-by-
220 environment interactions). Thus, despite strong temperature effects on average MIG, environmental
221 temperature appeared to have negligible effects on the relationship between genotype and phenotype.

222 **(c) Migrant probability and body size in late summer and show high correlations**

223 To test whether phenotypic body length in late summer contributes to MIG liability, we expanded the
224 reduced univariate model by adding phenotypic body length as a temperature-specific continuous
225 predictor ("length"). We expected that if length contributes to (or even constitutes) the liability, it would
226 positively covary with MIG. Furthermore, we expected that if genetic effects contribute positively to this
227 covariance, the explicit modelling of this covariance as slope would reduce MIG heritability. Length was
228 indeed a strong phenotypic predictor of MIG (**figure 2**). Controlled for length, temperature effects on MIG
229 became dependent on the size at which the contrast was made because the slope between MIG and
230 length was estimated as steeper in the warm than the cold environment (liability scale contrast: -1.18, -

231 1.85--0.49; **figure 2**). Estimated at the overall geometric mean length, MIG was higher in the cold than the
 232 warm environment (liability scale contrast: -1.18, -1.8--0.47).

233 Controlling for length also reduced heritability similarly in both environments (i.e., the heritability
 234 difference between environments was similar regardless of whether length was controlled for; **figure 3**).
 235 Heritability became non-significant in the warm but not the cold environment ($h^2_{\text{Cold}} = 0.26$ 0.11-0.42;
 236 $h^2_{\text{Warm}} = 0.12$, 0.00-0.28; **figure 3 a, b**), and controlling for length also reduced the between-environment
 237 genetic correlation (**figure 3 c**). Thus, phenotypic length in late summer explained a considerable share of
 238 MIG heritability (a phenotype expressed eight months later) in both temperature environments (52 and
 239 71%, respectively).

240 Having established that MIG covaried with phenotypic length, we quantified the phenotypic covariance
 241 and the relative contributions of environmental and genetic effects by fitting a bivariate model for MIG
 242 and body length in late summer as a second modelled response (LEN). Bivariate model estimates for
 243 average MIG in each temperature met those by the univariate model not controlling for phenotypic length
 244 ($\text{MIG}_{\text{Cold}} = 0.42$, 0.28-0.57; $\text{MIG}_{\text{Warm}} = 0.76$, 0.63-0.88). Estimates for average LEN in each temperature
 245 indicated strong temperature effects, whereby average LEN was 1.3 (1.1-1.5) phenotypic standard
 246 deviations (equating to 25%, 20-30%) larger in the warm than the cold environment. Back-transformed
 247 average LEN was 8.2 cm (7.8-8.7) in the warm and 6.6 cm (6.3-6.9) in the cold environment. For LEN, we
 248 estimated, unlikely for MIG, common environmental (tank) effects in the cold, but not the warm,
 249 environment that accounted for 9 % of the phenotypic variance (c^2 ; electronic supplementary material,
 250 **figure S2**).

251 Bivariate model estimates for MIG heritability were somewhat higher than by the univariate model (MIG;
 252 $h^2_{\text{Cold}} = 0.60$, 0.38-0.78, $h^2_{\text{Warm}} = 0.48$, 0.26-0.69; **figure 3**; electronic supplementary material, **figure S2**)
 253 and we estimated also high LEN heritabilities, which was surprisingly similar to the MIG heritabilities (LEN;
 254 $h^2_{\text{Cold}} = 0.57$, 0.38-0.74; $h^2_{\text{Warm}} = 0.52$, 0.30-0.73; **figure 3 a, b**). Using alternative prior specifications for the
 255 variances of MIG, heritabilities for both traits were estimated somewhat lower but their difference
 256 remained similar (**figure S3**). The between-temperature genetic correlations ($R_{\text{G Cold, Warm}}$) were also high
 257 and very similar for both MIG and LEN and their credible intervals were very close to unity (all $R_{\text{G Cold, Warm}} =$
 258 0.92; **figure 3 a, b**). These remarkably similar genetic correlations estimates differed only marginally with
 259 alternative prior specifications for the genetic covariances (electronic supplementary material, **figure S4**).
 260 Translated to the probability scale, we found MIG heritabilities to be lower and less similar between
 261 environments than on the liability scale (MIG; $h^2_{\text{Cold}} = 0.37$, 0.23-0.49; $h^2_{\text{Warm}} = 0.26$, 0.13-0.38), but the
 262 95% credible interval of the contrast on the probability scale also encompassed zero (MIG; $h^2_{\text{Cold}} - h^2_{\text{Warm}}$

263 = 0.12, -0.06-0.28). Thus, heritability and between-temperature genetic correlation estimates were similar
 264 for MIG (on the liability scale) and the liability proxy trait LEN.

265 As expected from univariate modelling, we detected high positive phenotypic correlations (R_P) between
 266 MIG and LEN in both environments ($R_{P_{Cold}} = 0.77$, 0.69-0.85; $R_{P_{Warm}} = 0.73$, 0.62-0.83; **figure 3 a, b**). The
 267 phenotypic correlations within temperature environments underlaid both genetic and environmental
 268 correlations, whereby the former exceeded the latter; the between-trait genetic correlation (R_G) was with
 269 0.92 consistently high in both environments and thereby equal to the between-temperature genetic
 270 correlation estimates for each trait. Between-trait environmental correlation estimates (R_E), for both
 271 temperature environments were somewhat lower ($R_{E_{Cold}} = 0.60$, 0.39-0.79; $R_{E_{Warm}} = 0.57$, 0.31-0.79; **figure**
 272 **3 a, b**), and even lower when using alternative prior specifications for the residual covariances (electronic
 273 supplementary material, **figure S4**). Relative contributions (weights) to the phenotypic correlation were
 274 either stronger or equal for genetic than environmental effects, whereby weights equate to the relative
 275 importance of the genetic and non-genetic components, respectively [46]. Specifically, the weights for
 276 genetic effects were 0.58 in the cold and 0.50 in the warm environment. As detected for MIG in the
 277 univariate model, no difference between temperatures was detected for variance parameters of either
 278 MIG or LEN (**figure 3**). Thus, correlation estimates between MIG and LEN were higher at the genetic than
 279 environmental level and both traits showed similar heritability and high correlation estimates across
 280 environmental temperatures, even though average expression of each trait differed considerably
 281 between temperature environments.

282 4. Discussion

283 The basis for variation in the developmental switches that result in categorical phenotypes remains a
 284 major research topic in several fields, including biology and medicine. The control of such categorical
 285 phenotypic variation may occur via (one or many) thresholds for the quantitative expression of a, usually
 286 unknown, liability that underlies environmental and genetic effects [16-18]. In this study, we investigated
 287 the developmental switch that determines whether Atlantic salmon express the migration phenotype
 288 associated with either a sea migration at one year of age (migrant) or with remaining in freshwater for
 289 longer (resident), and tested for association with its presumed liability proxy, body size in late summer.
 290 We estimated relatively high heritabilities for both migrant probability liability and its liability proxy trait,
 291 and high environmental, genetic, and phenotypic correlations between the two traits in two seasonal
 292 temperature environments. Furthermore, we estimated high between-environment genetic correlations
 293 for each trait. We, however, failed to detect male maturation, maternal, and feed restriction effects on
 294 migration phenotypes. Altogether, the results widen our understanding of genetic and environmental
 295 importance in the developmental transition towards age-specific migration phenotypes, as well as their

296 strong thermal stability, and have implications for ecological and evolutionary subjects and future studies,
297 which we elaborate on below.

298 **(a) Absence of male maturation and maternal effects on migration probability**

299 Our results strengthen previous findings, under both wild and culture conditions, that sexual maturation
300 of males in autumn is not inhibiting the migrant phenotype expression the following spring [36, 47-49].
301 Thus, the view that maturation and smoltification are conflicting processes [50], and which is also under
302 debate for the congeneric brown trout [15], may not hold generally in Atlantic salmon. We also rejected
303 the hypothesis that maternal effects affect migration phenotypes [such as suggested in 15], agreeing with
304 previous results in rainbow trout [51].

305 **(b) The relationship between migrant probability and body length**

306 Our results on Atlantic salmon strongly support a long-suspected, but often questioned, joint-governance
307 of body size in late summer and migration phenotype expression the following spring [9, 24]. We
308 demonstrate that, for a given temperature environment, this joint governance is stronger for genetic than
309 environmental effects, and that the genetic effects were extremely stable between two temperature
310 environments (**figure 3**). The results, thereby, provide quantitative primers for future modelling relying
311 on environmental vs. genetic relationships among life-history traits across environmental temperatures
312 [e.g., 4, 26, 33]. Given that the tested temperature difference of 2°C aligns with global warming scenarios
313 [52], modelling may encompass evolution under current global warming. Specifically, the combined high
314 between-trait and between-environment genetic correlation estimates suggest strong co-evolution of
315 migration phenotypes and body size across temperature environments, which may, depending on
316 ecological settings [9], be an advantage or a constraint.

317 We estimated high and similar heritability estimates for migrant probability and body size and high genetic
318 correlations, but only moderate environmental (residual) correlations (**figure 3**). Thus, the association
319 between traits appeared stronger for the genetic than the realised phenotypic values, which suggests that
320 genetic growth potential is also a proxy for migrant probability and not only size during a sensitive period.
321 An explanation could be that unknown genetically determined key components, such as growth hormone
322 levels or its receptor densities, may commonly control body size and migrant probability but need not
323 correlate tightly with phenotypic size during the sensitive period. Identifying such components may
324 explain how a scaling of the *genetic* growth potential occurs relative to actual body size, which varies
325 *environmentally* (such as by our temperature environments). Answers to this question may require more
326 detailed genetic mapping and in-depth investigations on the actual biological mechanism involved.

327 Given the time lag between the presumed sensitive period (in late summer) and expression of the
328 migration phenotype (the following spring), there may be time to catch up on missed growth

opportunities after the sensitive period. Thus, if we assume that a liability threshold ensures viable migrant sizes, phenotypic size in late summer may be a poorer predictor of migrant size in spring than genetic growth potential. In support of this, individuals initially falling within the lower group of a bimodal distribution in late summer or autumn, assumed to represent prospective residents, may still grow rapidly during winter and become migrants the following spring, both in captivity and in the wild [47, 53, 54]. As an alternative, the sensitive period may last longer than suspected. In that case, individuals with a high genetic potential for, but low realised, growth until late summer may still realise their genetic potential thereafter, thereby reaching a *phenotypic* size threshold later and initiate the transition later. Given that a growth spurt occurs in prospective migrants between late summer and winter as a consequence of the developmental transition [25, 55], it may be difficult to differentiate between “normal” and “spurt” growth to test between competing hypotheses. Nonetheless, the lack of effects on migration phenotypes by the feed restriction, applied after the presumed period, argues against the presence of an extended sensitive period (see below). Regardless of the sensitive period duration, our results suggest that individuals with relatively small body size in late summer may still become migrants if they possess a high genetic growth potential.

The presence of a higher genetic than environmental correlation between migrant probability and body size (**figure 3**) has several implications. For example, investigations on the physiological mechanisms of the developmental transition rely on body-size based migration-phenotype prediction for lethally sampling prospective migrants and residents prior to expression of the migration phenotype. Our results suggest that body-size based phenotype prediction may lead to conservative results because small migrants may get incorrectly assigned resident status. Such wrong assignments may blur differences of studied variables between true prospective migrants and residents. Another implication pertains to identifying the genetic basis for migration phenotypes via locus associations such as genome-wide association, or gene transcription studies, which may easily be confounded with identifying the highly polygenic basis for growth because of the high genetic between-trait correlation. To identify genes underlying also, or solely, the migration phenotype, a bivariate approach including size during the sensitive period may be necessary. Investigations on associations based on transcription levels during the sensitive period may also be promising, but pose logistical challenges due to the abovementioned phenotype prediction required for lethal sampling.

There are several inferential limitations to our estimates. Genetic parameter estimates often pertain to specific populations and conditions, but knowledge about their magnitudes may still provide useful information [56, 57]. Our estimates within and between temperature environments, may then be useful for future short-term predictions in response to selection under changing environmental conditions [58]. However, it is important to remember that heritabilities on the liability scale (and also heritabilities for

liability proxy traits) do not relate linearly to the probability scale across many factors, including the environmentally governed - overall probability [18]. This effect was here exemplified by disparate heritability differences between temperature environments on the liability (more similar) vs. the proportional (less similar) scale. As recently discussed by de Villemereuil, Schielzeth [45], predicting the responses to selection for migration phenotypes may follow standard assumptions if based on liability-scale heritability, but less so on proportional-scale heritability.

Pertaining to correlation estimates, Cheverud [59] proposed that many genetic, but not phenotypic, correlations estimates may be inflated, especially under sample size limitations and when heritabilities are low [59, 60]. In comparison to studies assessed by Cheverud [59], our effective sample sizes - the product of number of families and geometric mean heritability - are at the lower end where Cheverud [59] suspected upwards bias. However, Bayesian heritability estimates for length matched those by REML that should yield reliable estimates [43], and which supports the presence of moderate to high heritabilities. For migrant probability estimates, we rely on comparisons with previous estimates. To our knowledge, the only study in Atlantic salmon estimated a small heritability on the proportional scale (0.16 ± 0.05) at a lower migrant rate (0.18) than in the present study [55]. In that study migrants were defined as exceeding a particular size threshold in spring [55], altogether making a comparison difficult. In other salmonids, similarly high estimates for migrant probability liability as presented here exist, such as 0.61 in cultured rainbow trout [51] and 0.52-0.56 in wild brook charr [61]. Thus, migrant probability liability in salmonids may exhibit high heritabilities, which lends some support to the correlation estimates. Furthermore, previous genetic correlations estimates in rainbow trout between body length at age 12 and 15 months and migrant probability one year later, under much larger sample sizes, also were relatively high and exceeded the phenotypic correlation [51]. Thus, the detected stronger genetic than environmental relationship may not, at least entirely, be a statistical artefact.

(c) Temperature effects on migration phenotypes: liability vs. threshold variation

Threshold model interpretations affect biological inferences, including inferences about the here-detected temperature difference. Using the threshold model (or its variants) it is generally not possible to discern variation for thresholds from liability [41, 62]. This statistical uncertainty extends to general definitions. As a hypothetical example, it may be unclear whether circulating hormones and their receptors should be regarded as contributing commonly to liability variation, or the latter to threshold variation [as suggested in 24]. Here, we did not adopt either view when interpreting our results, but discuss all possibilities.

A possible interpretation for the detected temperature difference for average migrant probability (not controlled for length) could be that temperature affects body length positively. This more rapid growth in

the warm environment would then increase migrant probability because more individuals exceed a size threshold during a sensitive period. Assuming equal thresholds between temperatures, this idea leads to the expectation that controlling for size results in equal migrant rates. However, length-controlled migrant probability was higher in the cold environment (**figure 2**). This result could then be interpreted as the presence of a lower size threshold in the cold environment, supported by the high genetic correlations between environments as predicted in [62]. Nonetheless, because the exact sensitive period remains unknown, the length measured at a presumed sensitive period may be different than the length expressed at the actual sensitive period. This bias between approximated and actual liability proxy length may be larger in the warm than the cold environment simply because growth proceeds more rapidly in the warm environment. Thus, subtle methodological bias can create statistically different migrant rates at a standardised length that mimic the presence of different environmental (or in other cases: population) thresholds.

To not confound liability proxy bias with differences for environmental (or population) thresholds it would be necessary to base inferences on the true liability, or, when information is lacking, on the liability proxy but as expressed at the true sensitive period. This leads to the question how close we got to the true period. The developmental transition is inducible by shortening daylength from 24 to 12 h for 6 weeks [63] and may occur when daylength is shortened to between 15 and 12 h [30]. For our experimental setting, a 6-week period after daylengths got shorter, or daylengths between 15 and 12 h, occurred shortly before, or at the beginning of, the size recording period at the end of the summer and the middle of the feed restriction, respectively (**figure 1 c**). Because the feed restriction reduced specific growth rates by about 50% but not migrant probability, it appears likely that the sensitive period preceded the feed restriction, as similarly inferred for amago salmon [64]. Future experiments destined to identify the sensitive period - a required prior for sound investigations on the physiological or genetic mechanisms determining the developmental transition - may thus be worthwhile if covering the entire summer.

Data accessibility

Underlying data and R-scripts are available on the Dryad Digital Repository: https://datadryad.org/stash/share/clrV_-cHbjDel14IKMf9BnLUZiqublsJeny0seSzvX8

Authors' contributions

P.D. and C.P. conceived the study, C.P. and J.E. contributed materials, C.P. coordinated genotyping, P.D. and N.P. performed experiments and recorded data, P.D. analysed data and wrote the manuscript. All authors critically revised the article and contributed to its final version.

427 **Competing interests**

428 We declare no competing interests.

429 **Funding**

430 This work was supported by the Academy of Finland (project numbers 284941, 286334, 314254, and
431 314255), the European Research Council under the European Articles Union's Horizon 2020 research and
432 innovation program (grant agreement number 742312), and the University of Helsinki.

433 **Acknowledgements**

434 We thank S. Andrew, O. Andersson, T. Aykanat, Y. Czorlich, A. House, J. Kurko, M. Lindqvist, N. Lorenzen,
435 O. Mehtälä, K. Mobley, J. Moustakas-Verho, O. Ovaskainen, S. Papakostas, N. Parre, A. Ruokolainen, V.
436 Pritchard, K. Salminen, M. Sinclair-Waters, S. Tillanen, J.-P. Verta, K. Zueva, for help related to gamete
437 stripping, sample processing, tagging, genotyping, phenotyping, or fish husbandry and the staff at
438 Taivalkoski hatchery for help during spawning.

439

440 **Tables**

441 **Table 1.** Model mean posterior estimates (on the liability scale), lower and upper 95% credible intervals,
442 and number of effective samples (N_{eff}) for the initial univariate generalized animal model on migration
443 phenotype binaries of 663 Atlantic salmon individuals from 32 half-sib families. Variances were modelled
444 as either diagonal (tanks, residuals) or unstructured (dams, animals) covariance matrices for the two
445 temperature environments. Residual variance was fixed to 1 in each environment, resulting in scaling of all
446 components relative to the residual variance. Effects or variances different from zero (i.e., credible interval
447 not including zero) are in bold.

term	mean	lower	upper	N_{eff}
mean effects				
model intercept	-0.467	-1.325	0.423	15392
temperature (cold - warm)	1.942	0.988	2.920	13242
feed (full - restricted)	-0.010	-0.687	0.677	14000
maturation (immature - mature)	1.147	-0.892	3.161	14669
feed:temperature	-0.514	-1.607	0.537	14000
feed:maturation	-1.130	-3.536	1.151	14000
variance effects				
tank cold	0.127	0.000	0.455	13432
tank warm	0.166	0.000	0.608	14000
dam cold	0.182	0.000	0.691	13359
dam cold,warm	0.046	-0.139	0.319	13601
dam warm	0.188	0.000	0.694	14000
animal cold	2.677	0.641	4.766	14000
animal cold,warm	1.709	0.496	3.145	13430
animal warm	1.709	0.319	3.322	14000

448

References

1. Burghardt L.T., Metcalf C.J., Wilczek A.M., Schmitt J., Donohue K. 2015 Modeling the influence of genetic and environmental variation on the expression of plant life cycles across landscapes. *Am Nat* **185**(2), 212-227. (doi:10.1086/679439).
2. Huijser P., Schmid M. 2011 The control of developmental phase transitions in plants. *Development* **138**(19), 4117-4129. (doi:10.1242/dev.063511).
3. Fusco G., Minelli A. 2010 Phenotypic plasticity in development and evolution: facts and concepts. Introduction. *Philos Trans R Soc Lond B Biol Sci* **365**(1540), 547-556. (doi:10.1098/rstb.2009.0267).
4. Marschall E.A., Quinn T.P., Roff D.A., Hutchings J.A., Metcalfe N.B., Bakke T.A., Saunders R.L., Poff N.L. 1998 A framework for understanding Atlantic salmon (*Salmo salar*) life history. *Can J Fish Aquat Sci* **55**(S1), 48-58. (doi:10.1139/cjfas-55-S1-48).
5. Fox R.J., Donelson J.M., Schunter C., Ravasi T., Gaitan-Espitia J.D. 2019 Beyond buying time: the role of plasticity in phenotypic adaptation to rapid environmental change. *Philos Trans R Soc Lond B Biol Sci* **374**(1768), 20180174. (doi:10.1098/rstb.2018.0174).
6. Settele J., Scholes R., Betts R., Bunn S., Leadley P., Nepstad D., Overpeck J.T., Taboada M.A. 2014 Terrestrial and inland water systems. In *Climate Change 2014: Impacts, Adaptation, and Vulnerability Part A: Global and Sectoral Aspects* (eds. Field C.B., Barros V.R., Dokken D.J., Mach K.J., Mastrandrea M.D., Bilir T.E., Chatterjee M., Ebi K.L., Estrada Y.O., Genova R.C., et al.), pp. 271-359. Cambridge, Cambridge University Press.
7. Lack D. 1943 The problem with partial migration. *Br Birds* **37**(122-130).
8. Chapman B.B., Brönmark C., Nilsson J.-Å., Hansson L.-A. 2011 The ecology and evolution of partial migration. *Oikos* **120**(12), 1764-1775. (doi:10.1111/j.1600-0706.2011.20131.x).
9. Jonsson B., Jonsson N. 1993 Partial migration - niche shift versus sexual-maturation in fishes. *Rev Fish Biol Fish* **3**(4), 348-365. (doi:10.1007/Bf00043384).
10. Pulido F. 2011 Evolutionary genetics of partial migration - the threshold model of migration revis(it)ed. *Oikos* **120**(12), 1776-1783. (doi:10.1111/j.1600-0706.2011.19844.x).
11. Pearse D.E., Barson N.J., Nome T., Gao G., Campbell M.A., Abadia-Cardoso A., Anderson E.C., Rundio D.E., Williams T.H., Naish K.A., et al. 2019 Sex-dependent dominance maintains migration supergene in rainbow trout. *Nat Ecol Evol* **3**(12), 1731-1742. (doi:10.1038/s41559-019-1044-6).
12. Kelson S.J., Miller M.R., Thompson T.Q., O'Rourke S.M., Carlson S.M. 2019 Do genomics and sex predict migration in a partially migratory salmonid fish, *Oncorhynchus mykiss*? *Can J Fish Aquat Sci* **76**(11), 2080-2088. (doi:10.1139/cjfas-2018-0394).


- 481 13. Weinstein S.Y., Thrower F.P., Nichols K.M., Hale M.C. 2019 A large-scale chromosomal inversion
482 is not associated with life history development in rainbow trout from Southeast Alaska. *PLoS One* **14**(9),
483 e0223018. (doi:10.1371/journal.pone.0223018).
- 484 14. Chapman B.B., Hulthen K., Brodersen J., Nilsson P.A., Skov C., Hansson L.A., Bronmark C. 2012
485 Partial migration in fishes: causes and consequences. *J Fish Biol* **81**(2), 456-478. (doi:10.1111/j.1095-
486 8649.2012.03342.x).
- 487 15. Ferguson A., Reed T.E., Cross T.F., McGinnity P., Prodohl P.A. 2019 Anadromy, potamodromy and
488 residency in brown trout *Salmo trutta*: the role of genes and the environment. *J Fish Biol* **95**(3), 692-718.
489 (doi:10.1111/jfb.14005).
- 490 16. Falconer D.S. 1965 The inheritance of liability to certain diseases, estimated from the incidence
491 among relatives. *Ann Hum Genet* **29**(1), 51-76. (doi:10.1111/j.1469-1809.1965.tb00500.x).
- 492 17. Wright S. 1934 An analysis of variability in number of digits in an inbred strain of guinea pigs.
493 *Genetics* **19**(6), 506-536.
- 494 18. Dempster E.R., Lerner I.M. 1950 Heritability of threshold characters. *Genetics* **35**(2), 212-236.
- 495 19. Hoar W.S. 1988 The Physiology of Smolting Salmonids. In *FISH PHYSIOLOGY The Physiology of*
496 *Developing Fish: Viviparity and Posthatching Juveniles* (eds. Hoar W.S., Randall D.J.), pp. 275-343. San
497 Diego, California, Academic Press.
- 498 20. Salminen M., Kuikka S., Erkamo E. 1995 Annual variability in survival of sea-ranched Baltic salmon,
499 *Salmo salar* L: significance of smolt size and marine conditions. *Fish Manage Ecol* **2**(3), 171-184.
500 (doi:10.1111/j.1365-2400.1995.tb00110.x).
- 501 21. Thorstad E.B., Whoriskey F., Uglem I., Moore A., Rikardsen A.H., Finstad B. 2012 A critical life stage
502 of the Atlantic salmon *Salmo salar*: behaviour and survival during the smolt and initial post-smolt
503 migration. *J Fish Biol* **81**(2), 500-542. (doi:10.1111/j.1095-8649.2012.03370.x).
- 504 22. Chaput G., Carr J., Daniels J., Tinker S., Jonsen I., Whoriskey F. 2018 Atlantic salmon (*Salmo salar*)
505 smolt and early post-smolt migration and survival inferred from multi-year and multi-stock acoustic
506 telemetry studies in the Gulf of St. Lawrence, northwest Atlantic. *ICES J Mar Sci.*
507 (doi:10.1093/icesjms/fsy156).
- 508 23. Gregory S.D., Ibbotson A.T., Riley W.D., Nevoux M., Lauridsen R.B., Russell I.C., Britton J.R.,
509 Gillingham P.K., Simmons O.M., Rivot E., et al. 2019 Atlantic salmon return rate increases with smolt
510 length. *ICES J Mar Sci.* (doi:10.1093/icesjms/fsz066).
- 511 24. Dodson J.J., Aubin-Horth N., Theriault V., Paez D.J. 2013 The evolutionary ecology of alternative
512 migratory tactics in salmonid fishes. *Biol Rev Camb Philos Soc* **88**(3), 602-625. (doi:10.1111/brv.12019).
- 513 25. Thorpe J.E. 1977 Bimodal distribution of length of juvenile Atlantic salmon (*Salmo salar* L.) under
514 artificial rearing conditions. *J Fish Biol* **11**(2), 175-184. (doi:10.1111/j.1095-8649.1977.tb04111.x).


- 515 26. Mangel M., Satterthwaite W.H. 2008 Combining proximate and ultimate approaches to
516 understand life history variation in salmonids with application to fisheries, conservation, and aquaculture.
517 *Bull Mar Sci* **83**(1), 107-130.
- 518 27. Björnsson B.T., Stefansson S.O., McCormick S.D. 2011 Environmental endocrinology of salmon
519 smoltification. *Gen Comp Endocrinol* **170**(2), 290-298. (doi:10.1016/j.ygcen.2010.07.003).
- 520 28. Kristinsson J.B., Saunders R.L., Wiggs A.J. 1985 Growth dynamics during the development of
521 bimodal length-frequency distribution in juvenile Atlantic salmon (*Salmo salar* L.). *Aquaculture* **45**(1-4), 1-
522 20. (doi:10.1016/0044-8486(85)90254-6).
- 523 29. Metcalfe N.B., Huntingford F.A., Graham W.D., Thorpe J.E. 1989 Early social status and the
524 development of life-history strategies in Atlantic salmon. *Proc R Soc B* **236**(1282), 7-19.
525 (doi:10.1098/rspb.1989.0009).
- 526 30. Skilbrei O.T., Hansen T., Stefansson S.O. 1997 Effects of decreases in photoperiod on growth and
527 bimodality in Atlantic salmon *Salmo salar* L. *Aquacult Res* **28**(1), 43-49. (doi:10.1046/j.1365-
528 2109.1997.00827.x).
- 529 31. Wright P.J., Metcalfe N.B., Thorpe J.E. 1990 Otolith and somatic growth rates in Atlantic salmon
530 parr, *Salmo salar* L: evidence against coupling. *J Fish Biol* **36**(2), 241-249. (doi:10.1111/j.1095-
531 8649.1990.tb05599.x).
- 532 32. Bailey J.K., Saunders R.L., Buzeta M.I. 1980 Influence of parental smolt age and sea age on growth
533 and smolting of hatchery-reared Atlantic salmon (*Salmo salar*). *Can J Fish Aquat Sci* **37**(9), 1379-1386.
534 (doi:10.1139/f80-177).
- 535 33. de Zoeten T., Pulido F. 2020 How migratory populations become resident. *Proc R Soc B* **287**(1923),
536 20193011. (doi:10.1098/rspb.2019.3011).
- 537 34. Karppinen P., Jounela P., Huusko R., Erkinaro J. 2014 Effects of release timing on migration
538 behaviour and survival of hatchery-reared Atlantic salmon smolts in a regulated river. *Ecol Freshwat Fish*
539 **23**(3), 438-452. (doi:10.1111/eff.12097).
- 540 35. Verta J.-P., Debes P.V., Piavchenko N., Ruokolainen A., Ovaskainen O., Moustakas-Verho J.E.,
541 Tillanen S., Parre N., Aykanat T., Erkinaro J., et al. 2019 Regulatory divergence in *vgll3* underlies variation
542 in age at maturity in male Atlantic salmon. *bioRxiv*. (doi:10.1101/777300).
- 543 36. Bagliniere J.L., Maisse G. 1985 Precocious maturation and smoltification in wild Atlantic salmon
544 in the Armorican massif, France. *Aquaculture* **45**(1-4), 249-263. (doi:10.1016/0044-8486(85)90274-1).
- 545 37. Aykanat T., Lindqvist M., Pritchard V.L., Primmer C.R. 2016 From population genomics to
546 conservation and management: a workflow for targeted analysis of markers identified using genome-wide
547 approaches in Atlantic salmon *Salmo salar*. *J Fish Biol* **89**(6), 2658-2679. (doi:10.1111/jfb.13149).
- 548 38. Jones O.R., Wang J. 2010 COLONY: a program for parentage and sibship inference from multilocus
549 genotype data. *Mol Ecol Resour* **10**(3), 551-555. (doi:10.1111/j.1755-0998.2009.02787.x).

- 550 39. Anderson E.C. 2010 Computational algorithms and user-friendly software for parentage-based
551 tagging of Pacific salmonids. Final report submitted to the Pacific Salmon Commission's Chinook Technical
552 Committee (US Section). (p. 46.
- 553 40. Henderson C.R. 1973 Sire evaluation and genetic trends. *J Anim Sci* **1973**(Symposium), 10-41.
554 (doi:/1973.1973Symposium10x).
- 555 41. Hadfield J.D., O'Hara R.B. 2015 Increasing the efficiency of MCMC for hierarchical phylogenetic
556 models of categorical traits using reduced mixed models. *Methods Ecol Evol* **6**(6), 706-714.
557 (doi:10.1111/2041-210x.12354).
- 558 42. Hadfield J.D. 2010 MCMC methods for multi-response generalized linear mixed models: the
559 MCMCglmm R package. *J Stat Soft* **33**(2), 1-22. (doi:10.18637/jss.v033.i02).
- 560 43. de Villemereuil P., Gimenez O., Doligez B., Freckleton R. 2013 Comparing parent-offspring
561 regression with frequentist and Bayesian animal models to estimate heritability in wild populations: a
562 simulation study for Gaussian and binary traits. *Methods Ecol Evol* **4**(3), 260-275. (doi:10.1111/2041-
563 210x.12011).
- 564 44. Brooks S.P., Gelman A. 1998 General methods for monitoring convergence of iterative
565 simulations. *J Comput Graph Stat* **7**(4), 434-455. (doi:10.1080/10618600.1998.10474787).
- 566 45. de Villemereuil P., Schielzeth H., Nakagawa S., Morrissey M. 2016 General methods for
567 evolutionary quantitative genetic inference from generalized mixed models. *Genetics* **204**(3), 1281-1294.
568 (doi:10.1534/genetics.115.186536).
- 569 46. Searle S.R. 1961 Phenotypic, genetic and environmental correlations. *Biometrics* **17**(3), 474-480.
570 (doi:10.2307/2527838).
- 571 47. Duston J., Saunders R.L. 1997 Life histories of Atlantic salmon altered by winter temperature and
572 summer rearing in fresh-or sea-water. *Environ Biol Fishes* **50**(2), 149-166.
573 (doi:10.1023/A:1007373016067).
- 574 48. Orton J.H., Jones J.W., King G.M. 1938 The male sexual stage in salmon parr (*Salmo salar* L. juv.).
575 *Proc R Soc B* **125**(838), 103-114. (doi:10.1098/rspb.1938.0015).
- 576 49. Jonsson N., Jonsson B., Hansen L.P. 1998 Long-term study of the ecology of wild Atlantic salmon
577 smolts in a small Norwegian river. *J Fish Biol* **52**(3), 638-650. (doi:10.1111/j.1095-8649.1998.tb02023.x).
- 578 50. Thorpe J.E. 1986 Age at first maturity in Atlantic salmon, *Salmo salar*: freshwater period influences
579 and conflicts with smolting. In *Salmonid age at maturity* (ed. Meerburg D.J.), pp. 7-14. Ottawa,
580 Department of Fisheries and Oceans.
- 581 51. Hecht B.C., Hard J.J., Thrower F.P., Nichols K.M. 2015 Quantitative genetics of migration-related
582 traits in rainbow and steelhead trout. *G3 (Bethesda)* **5**(5), 873-889. (doi:10.1534/g3.114.016469).
- 583 52. IPCC. 2014 Future Climate Changes, Risk and Impacts. In *Climate Change 2014: Synthesis Report*
584 *Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel*

- 585 on *Climate Change* (eds. Core Writing Team, Pachauri R.K., Meyer L.A.), pp. 56-74. Geneva, Switzerland,
586 IPCC.
- 587 53. Zydlewski J., O'Malley A., Cox O., Ruksznis P., Trial J.G. 2014 Growth and smolting in lower-mode
588 Atlantic salmon stocked into the Penobscot River, Maine. *N Am J Fish Manage* **34**(1), 147-158.
589 (doi:10.1080/02755947.2013.866996).
- 590 54. Whitesel T.A. 1993 Comparison of juvenile Atlantic salmon (*Salmo salar*) reared in a hatchery and
591 introduced into a stream: a two-size-threshold model for smoltification. In *Production of Juvenile Atlantic*
592 *salmon, Salmo salar, in Natural Waters* (eds. Gibson R.J., Cutting R.E.), pp. 239-247. Ottawa, Canada,
593 National Research Council Canada and Department of Fisheries and Oceans.
- 594 55. Refstie T., Steine T.A., Gjerdem T. 1977 Selection experiments with salmon. II. Proportion of
595 Atlantic salmon smoltifying at 1 year of age. *Aquaculture* **10**(3), 231-242. (doi:10.1016/0044-
596 8486(77)90004-7).
- 597 56. Hill W.G. 2010 Understanding and using quantitative genetic variation. *Philos Trans R Soc Lond B*
598 *Biol Sci* **365**(1537), 73-85. (doi:10.1098/rstb.2009.0203).
- 599 57. Visscher P.M., Hill W.G., Wray N.R. 2008 Heritability in the genomics era: concepts and
600 misconceptions. *Nat Rev Genet* **9**(4), 255-266. (doi:10.1038/nrg2322).
- 601 58. Falconer D.S. 1952 The problem of environment and selection. *Am Nat* **86**(830), 293-298.
602 (doi:10.1086/281736).
- 603 59. Cheverud J.M. 1988 A Comparison of genetic and phenotypic correlations. *Evolution* **42**(5), 958-
604 968. (doi:10.1111/j.1558-5646.1988.tb02514.x).
- 605 60. Robertson A. 1959 The sampling variance of the genetic correlation coefficient. *Biometrics* **15**(3),
606 469-485. (doi:10.2307/2527750).
- 607 61. Thériault V., Garant D., Bernatchez L., Dodson J.J. 2007 Heritability of life-history tactics and
608 genetic correlation with body size in a natural population of brook charr (*Salvelinus fontinalis*). *J Evol Biol*
609 **20**(6), 2266-2277. (doi:10.1111/j.1420-9101.2007.01417.x).
- 610 62. Roff D.A. 1996 The evolution of threshold traits in animals. *Q Rev Biol* **71**(1), 3-35.
611 (doi:10.1086/419266).
- 612 63. Björnsson B.T., Hemre G.I., Bjørnevik M., Hansen T. 2000 Photoperiod regulation of plasma
613 growth hormone levels during induced smoltification of underyearling Atlantic salmon. *Gen Comp*
614 *Endocrinol* **119**(1), 17-25. (doi:10.1006/gcen.2000.7439).
- 615 64. Kuwada T., Tokuhara T., Shimizu M., Yoshizaki G. 2015 Body size is the primary regulator affecting
616 commencement of smolting in amago salmon *Oncorhynchus masou ishikawae*. *Fish Sci* **82**(1), 59-71.
617 (doi:10.1007/s12562-015-0943-y).

618





Genetic growth potential, rather than phenotypic size, predicts migration phenotype in Atlantic salmon (doi: 10.1098/rspb.2020-0867)

Paul V Debes^{1,2*}, Nikolai Piavchenko¹, Jaakko Erkinaro³, Craig R Primmer^{1,4}

¹Organismal & Evolutionary Biology Research Program, University of Helsinki, Helsinki, 00014, Finland; ²Department of Aquaculture and Fish Biology, Hólar University College, Sauðárkrúkur, 551, Iceland; ³Natural Resources Institute Finland (Luke), Oulu, 90014, Finland; ⁴Institute of Biotechnology, University of Helsinki, Helsinki, 00014, Finland

*Correspondence: debes@holar.is

Contents:

- Table S1
- Description of the general linear mixed to test for feed restriction treatment effects
- Table S2
- Figure S1
- Description of the generalised linear mixed model to estimate male maturation rate
- Table S3
- Prior sensitivity analysis
- Figure S2
- Figure S3
- Figure S4
- References

Table S1. Model mean posterior estimates (on the liability scale), lower and upper 95% credible intervals, and number of effective samples (N_{eff}) for the reduced univariate generalized animal model on migrant phenotype binaries of 663 Atlantic salmon individuals from 32 half-sib families, including the effects of a locus with major effects on age at sexual maturation (*vgll3*, alleles: E = early, L = late maturation). Results for the binary response of migrant (1) or resident (0) are based on models not controlling for phenotypic length (model 1) or controlling for phenotypic length (model 2; length = length in late summer; mean centred and variance scaled).

Term	mean	lower	upper	N_{eff}
model 1				
model intercept	-0.338	-1.271	0.551	14516
temperature (cold - warm)	1.701	1.025	2.439	14564
Vgll3 (EE - EL)	-0.139	-0.780	0.512	14000
Vgll3 (EE - LL)	-0.276	-1.328	0.806	14000
model 2				
model intercept	1.287	0.616	1.913	7000
length (continuous)	2.946	2.435	3.507	7000
temperature (cold - warm)	-0.984	-1.660	-0.305	7000
temperature:length	-1.192	-1.864	-0.532	6317
Vgll3 (EE - EL)	0.164	-0.299	0.642	7000
Vgll3 (EE - LL)	0.213	-0.489	0.883	7000

Description of the general linear mixed to test for feed restriction treatment effects

To test for effect of the feed treatment (see methods for details) on growth rate, we fitted a general linear animal model under restricted residual maximum likelihood (REML) using asreml-R [1], to the (log) of length before and after the temporary feed restriction treatment. To estimate specific growth rate (i.e., proportional increase of length per unit time), we included date of measurement as a continuous covariate (date.integer; first measurement date taken as 0, days have a value of 1). We allowed the intercept (at the first measurement) and the growth rates to vary by temperature (cold, warm), feed restriction (full, restricted), and by migrant phenotype (resident, migrant). We also fitted all interactions among these terms. To account for the randomisation of treatments to tanks, we fitted random regression effects for tanks (i.e., a 2x2 covariance matrix for tank intercepts and date slopes with covariance between them). We accounted for the non-independence of, and among, individual data by including animal effects with additive genetic variance estimated via the inverse of the relationship matrix [2]. Using likelihood ratio tests between nested models, we determined that variances for animal effects ($X_1^2 = 17.79$, $P < 0.001$) and residuals ($X_1^2 = 17.79$, $P < 0.001$), but not tanks effects ($X_3^2 = 1.97$, $P = 0.580$), differed between temperature environments. We thus fitted these former two effect terms conditional on temperature environment, and with between-temperature covariance for the genetic effects. We found that the feed restriction had affected the specific growth rate of both residents and migrants and in both the cold and the warm environments (**table S2, figure S1**).

Table S2. ANOVA table for model terms of a univariate general animal model on body length phenotype in late summer of 663 Atlantic salmon individuals from 32 half-sib families. Intercept effects were estimated at the first measurement date (date.integer = 0). Statistical significance has been estimated based on *F*-tests with denominator degrees of freedom, DDF, approximated according to [3].

term	DF	DDF	<i>F</i>	<i>P</i>
model intercept	1	51.6	6162.0	< 0.001
migrant phenotype (resident, migrant)	1	518.8	448.4	< 0.001
temperature (cold - warm)	1	22.7	146.1	< 0.001
feed restriction (full - restricted)	1	11.8	7.6	0.018
date.integer (continuous)	1	11.3	904.1	< 0.001
migrant phenotype:temperature	1	206.0	8.8	0.003
migrant phenotype:feed restriction	1	550.9	0	0.968
temperature:feed restriction	1	14.7	0.1	0.782
migrant phenotype:date.integer	1	1302.0	430.9	< 0.001
temperature:date.integer	1	1302.0	9.7	0.002
feed restriction:date.integer	1	1302.0	122.0	< 0.001
migrant phenotype:temperature:feed restriction	1	407.8	0.3	0.607
migrant phenotype:temperature:date.integer	1	1302.0	8.1	0.004
migrant phenotype:feed restriction:date.integer	1	1302.0	38.3	< 0.001
temperature:feed restriction:date.integer	1	12.0	0.1	0.757
migrant phenotype:temperature:feed restriction:date.integer	1	409.3	10.8	0.001




Figure S1. Model predicted average size trajectories (*a, d*) and associated specific growth rates (SGR; *b, e*) for prospective migrants or residents that were either fully fed or temporarily restrictedly fed in either a cold (upper row) or a 2°C warmer (lower row) environment, and the corresponding contrasts for SGR between the feeding treatments (ΔSGR; *c, f*). Estimates refer to the model as in **table S2**.

Description of the generalised linear mixed model to estimate male maturation rate

To estimate male maturation rate, we fit a generalised linear animal model with probit-link function to maturation binaries recorded during spawning time. We only fitted one overall mean effect (intercept). To account for the randomisation to tanks, we fitted random effects for tanks and to account for the non-independence among individual data, we including animal effects with additive genetic variance estimated via the inverse of the relationship matrix [2]. We used priors following [4] and methods as described for univariate Bayesian models in the main manuscript. We found that among tank effect variance was absent or negligible but that additive genetic variance was present (although methodologically inflated; **table S2**) and predicted a marginal overall maturation rate of males in the warm environment of 0.19 (95% CI: 0.08-0.33; **table S2**).

Table S3. Model mean posterior estimates (on the liability scale), lower and upper 95% credible intervals, and number of effective samples (N_{eff}) for the univariate generalized animal model on the sexual maturation phenotype binaries (0 = immature, 1 = mature) during the first year of 114 Atlantic salmon individuals from 30 half-sib families. Estimates were obtained for only males (no female matured) and only in the warm environment (no maturation occurred in the cold environment). Residual variance was fixed to 1, resulting in scaling of all components relative to the fixed residual variance. Variances different from zero (approximated by credible interval not including zero) are in bold.

Term	mean	lower	upper	N_{eff}
mean terms				
model intercept	-1.647	-2.615	-0.758	10000
variance terms				
Tank (V_C)	0.417	0.000	1.449	10210
Animal (V_A)*	2.127	0.600	3.828	10000

*Estimate is inflated because it is based on the coefficients of relatedness among individuals, but genotypes for the locus with major effects on maturation rate (*vgll3*) did not vary within families (leading to an absence of the expecting Mendelian sampling variance within families) due to the breeding design of using only *vgll3* homozygous parents as reported in [5].

Prior sensitivity analysis

We also assessed how prior specifications influenced the results. We focussed on the binary trait (MIG) because Bayesian heritability estimates for LEN closely matched estimates by residual maximum likelihood (REML), although the genetic correlation between environments was higher under REML (**figure S2**), whereby REML is the recommend method for animal models of continuous responses [4]. We compared the bivariate results (responses: MIG, LEN; including GxE) with univariate results (MIG per environment) obtained with recommended priors for binary animal models (following a X_1^2 distribution) [4]. We also separately assessed the effects, relative to the univariate models, of either including GxE for MIG or also fitting LEN per temperature environment. For the bivariate model, we varied prior specifications that resulted in different prior distributions for the variances and covariances. For the variances, we varied the parameter expansion variance scale by up to two magnitudes higher and lower than used (100, 10, 1, 0.1, 0.01), resulting in increasing prior densities for the proportions of the phenotypic variances towards one and zero, respectively. For the covariances, and thus correlations, we specified prior distributions that were either relatively flat for correlations (in *MCMCglmm*: nu = dimension of C, G, or R + 1) or showed higher densities towards -1 and 1 (in *MCMCglmm*: nu = dimension of C, G, or R). We found all investigated modelling variants to affect model estimates, but none strong enough to compromise our major inferences (**figures S2-S4**).




Figure S2. Comparison of the Bayesian model estimates for (co)variance components with increasing model complexity for the binary (MIG) and with REML estimates for the continuous trait (LEN). Heritability estimates for MIG increased in the following order: univariate single environment (UV) < univariate two environments (UV + GxE) < bivariate single environment (BV) < bivariate two environments (BV + GxE). Unfortunately, it is not clear whether model estimates in the absence of an explicitly modelled between-environment correlation (GxE) for the additive genetic effects should equal estimates in the absence of such modelled GxE. Furthermore, it is not clear whether model estimates for a single trait should equal those for estimates when a correlated second traits is modelled. Error bars show 95% credible intervals or approximate confidence intervals based on the delta method for REML estimates.




Figure S3 Comparison of the Bayesian model estimates for (co)variance components under different prior specifications for the parameter expansion variance scale (in *MCMCglmm* specified as “ $\alpha.V$ ”) for the genetic and common environmental covariance matrices of the binary trait (MIG). For MIG heritability (h^2) and phenotypic proportion of the common environmental variance (c^2), the priors specify an increasing density towards 0 ($\alpha.V$ specified as either 100, 10, 1, 0.1, or 0.01). The parameter expansion variance scale specification for the continuous trait (LEN) was kept constant ($\alpha.V = 1$). It could be noted that a smaller variance scale resulted in smaller heritability (h^2) and proportion of common environmental variance (c^2) of MIG when $\alpha.V$ fell < 1 . Similar effects on h^2 , but not c^2 , were also observed for LEN.




Figure S4. Comparison of the MCMC bivariate model estimates for (co)variance components varying the prior specification for the degree of believe parameter (ν) for combinations of the genetic (G, also applied for the environmental C) and residual (R) covariance matrices. The different priors result in either relatively flat distributions for the correlations (in *MCMCglmm*: ν = dimension [dim] of G or R + 1) or distributions with higher densities towards 0 and 1 (in *MCMCglmm*: ν = dimension [dim] of G or R). We noted an effect on the between-trait correlation for the common environmental effects within the cold environment ($R_{C_{MIG,LEN}}$; **a**), which is unsurprising because common environmental effects were inferred as absent for one trait (c^2_{MIG}) but present for the other (c^2_{LEN}), thus lacking data information on correlations. As a result, we

refrained from making inferences about $R_{C_{MIG,LEN}}$, and effects of $R_{C_{MIG,LEN}}$ on $R_{P_{MIG,LEN}}$ were marginal because of the low or absent c^2 . A second effect could be noted on the between-trait covariance for the residual environmental effects within both temperature environments ($R_{E_{MIG,LEN}}$; **a, b**). Given the prior knowledge of a suspected causal relationship between the two traits and for each trait between environments, but uncertainty whether this correlation is stronger at the environmental or genetic level, it is unclear which prior is more appropriate (i.e., whether assuming a between-trait correlation closer to zero or unity). However, this difference affects inferences only marginally and genetic correlation estimates appeared unaffected.

References

1. Butler D.G., Cullis B.R., Gilmour A.R., Gogel B.J. 2009 *Mixed models for S language environments ASReml-R reference manual*. Brisbane, Australia, Queensland Department of Primary Industries and Fisheries, NSW Department of Primary Industries; 144 p.
2. Henderson C.R. 1973 Sire evaluation and genetic trends. *J Anim Sci* **1973**(Symposium), 10-41. (doi:/1973.1973Symposium10x).
3. Kenward M.G., Roger J.H. 1997 Small sample inference for fixed effects from restricted maximum likelihood. *Biometrics* **53**(3), 983-997. (doi:10.2307/2533558).
4. de Villemereuil P., Gimenez O., Doligez B., Freckleton R. 2013 Comparing parent-offspring regression with frequentist and Bayesian animal models to estimate heritability in wild populations: a simulation study for Gaussian and binary traits. *Methods Ecol Evol* **4**(3), 260-275. (doi:10.1111/2041-210x.12011).
5. Verta J.-P., Debes P.V., Piavchenko N., Ruokolainen A., Ovaskainen O., Moustakas-Verho J.E., Tillanen S., Parre N., Aykanat T., Erkinaro J., et al. 2019 Regulatory divergence in *vgl/3* underlies variation in age at maturity in male Atlantic salmon. *bioRxiv*. (doi:10.1101/777300).